

Relative sensitivity of five Hawaiian coral species to high temperature under high-pCO₂ conditions

Keisha D. Bahr¹ · Paul L. Jokiel¹ · Ku'ulei S. Rodgers¹

Received: 22 January 2015 / Accepted: 13 January 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract Coral reef ecosystems are presently undergoing decline due to anthropogenic climate change. The chief detrimental factors are increased temperature and increased pCO₂. The purpose of this study was to evaluate the effect of these two stressors operating independently and in unison on the biological response of common Hawaiian reef corals. Manipulative experiments were performed using five species (*Porites compressa*, *Pocillopora damicornis*, *Fungia scutaria*, *Montipora capitata*, and *Leptastrea purpurea*) in a continuous-flow mesocosm system under natural sunlight conditions. Corals were grown together as a community under treatments of high temperature (2 °C above normal maximum summer temperature), high pCO₂ (twice present-day conditions), and with both factors acting in unison. Control corals were grown under present-day pCO₂ and at normal summer temperatures. *Leptastrea purpurea* proved to be an extremely hardy coral. No change in calcification or mortality occurred under treatments of high temperature, high pCO₂, or combined high temperature–high pCO₂. The remaining four species showed reduced calcification in the high-temperature treatment. Two species (*L. purpurea* and *M. capitata*) showed no response to increased pCO₂. Also, high pCO₂ ameliorated the negative effect of high temperature on the

calcification rates of *P. damicornis*. Mortality was driven primarily by high temperature, with a negative synergistic effect in *P. compressa* only in the high-pCO₂–high-temperature treatment. Results support the observation that biological response to temperature and pCO₂ elevation is highly species-specific, so generalizations based on response of a single species might not apply to a diverse and complex coral reef community.

Keywords Coral calcification · Ocean acidification · Interactive effects · Synergy · Antagonistic · Climate change

Introduction

Coral reefs have immense biological wealth and provide economic and environmental services (e.g., tourism, fisheries, wave protection, food production) for millions of people. However, reefs throughout the world are undergoing significant ecological change due to anthropogenic climate change (Hoegh-Guldberg 1999; Pandolfi et al. 2011). A major concern is the decrease in the pH of the ocean due to increasing uptake of anthropogenic atmospheric pCO₂, a process referred to as ocean acidification (OA) (Hoegh-Guldberg et al. 2007; Erez et al. 2011). Ocean acidity has increased by approximately 25 % (decrease of 0.1 pH units) since pre-industrial times (Lacis et al. 2010) and has resulted in severe negative effects on calcifying marine organisms (Smith and Buddemeier 1992; Kleypas et al. 2006; Jokiel et al. 2008; Erez et al. 2011; Gattuso and Hansson 2011; Kroeker et al. 2013). The calcification rate of reef-building corals is predicted to decrease by 40 % with increasing pCO₂ concentrations by the end of the century (Gattuso et al. 1999; Kleypas et al.

Communicated by Biology Editor Dr. Anastazia Banaszak

Electronic supplementary material The online version of this article (doi:10.1007/s00338-016-1405-4) contains supplementary material, which is available to authorized users.

✉ Keisha D. Bahr
kbahr@hawaii.edu

¹ Hawai'i Institute of Marine Biology, University of Hawai'i, P.O. Box 1346, Kaneohe, HI 96744, USA

1999; Langdon et al. 2000; Marubini et al. 2001, 2003; Müller et al. 2004; Langdon and Atkinson 2005; Kleypas and Langdon 2006; Hofmann et al. 2010; Erez et al. 2011). Experimental and field studies show that the impact of increased pCO₂ on calcification rates varies among species (Gattuso et al. 1998; Marubini et al. 2001, 2003; Reynaud et al. 2003; Edmunds et al. 2012; Comeau et al. 2013b). These variations in OA sensitivities may lead to differences in coral and algal species composition and therefore modify habitat complexity in the future (Fabricius et al. 2014; Comeau et al. 2015).

Along with OA, excessive ocean warming is predicted to negatively affect corals through increased thermal bleaching, reduced calcification, and increased coral mortality (Jokiel and Coles 1977, 1990; Glynn and D'Croz 1990; Lesser et al. 1990; Brown 1997; Erez et al. 2011). Prolonged exposure to elevated sea surface temperatures (SST) (+2 °C above upper lethal maximum) decreases photosynthesis and photosynthetic efficiency in the algal symbionts (Brown 1997); therefore, photosynthesis-to-respiration ratio decreases with increasing temperatures (Coles and Jokiel 1977). Calcification declines sharply above peak summer temperature with bleaching and eventual death of corals under future temperature scenarios (Hoegh-Guldberg et al. 2007; Buddemeier et al. 2008; Hoeke et al. 2011). The occurrence and severity of mass coral bleaching has increased dramatically over the last two decades with almost every reef region in the world suffering extensive coral stress and mortality (Jokiel and Brown 2004; Buddemeier et al. 2008; Bahr et al. 2015b). The isolated subtropical coral reef communities of Hawai'i, which contain high proportions of marine endemic species (>25 % in most taxa), have only recently begun to suffer from conditions that have ravaged coral reef communities in other areas of the world (Bahr et al. 2015b).

Increases in SST and pCO₂ operating independently have been shown to be detrimental to corals, but less is known about their effect when operating in unison. Therefore, it is essential to assess direct impacts of multiple stressors and determine whether the combine effects exceed (synergism) or are less than (antagonism) the sum of their individual effects. Moreover, the biological response of calcifying organisms to these climate change stressors may be species-specific, which may explain conflicting results regarding coral response to OA and multiple climate change stressors (Ban et al. 2014). The objectives of this study were to characterize the relative biological response (calcification and partial mortality) of Hawaiian reef-building corals under elevated temperatures and high pCO₂ and to describe the interactive effect (synergism or antagonism) when both factors operate in unison. Five abundant Hawaiian scleractinian corals—*Porites compressa*, *Montipora capitata*, *Fungia scutaria*, *Leptastrea purpurea*, and *Pocillopora*

damicornis—were studied. These corals show a wide range of skeletal morphologies, tissue structure, documented stress tolerance levels, and reproductive strategies (Bahr et al. 2015a). The goal of the present study was to evaluate the direct and interactive effects of elevated temperature and acidification on the biological response of these corals in order to improve our ability to forecast future changes in species composition for Hawaiian reefs under projected global climate change conditions.

Materials and methods

Experimental system

This research was conducted at University of Hawai'i, Hawai'i Institute of Marine Biology (HIMB) at Moku o Lo'e, Kāne'ohe Bay, Hawai'i. The experiment was conducted in an outdoor continuous-flow experimental facility that mimics the physical, chemical, and biological conditions on the adjacent reef flat under projected conditions of climate change (Jokiel et al. 2008, 2014a). Carbonate chemistry within the mesocosms was manipulated through direct bubbling of CO₂ (Jokiel et al. 2014a), and temperature was controlled using titanium heaters (Finnex, TH-800, 800 W) (Table 1).


The stressors of acidification and temperature were manipulated within 500-L mesocosms to test each factor independently and provide a pairwise comparison of these variables during the summer months of June through August 2013. The experimental mesocosms were located in full sunlight (Fig. 1; Table 2) and allowed for natural diurnal fluctuations in seawater chemistry under present and future climate change projections (Jokiel et al. 2014a) (Fig. 2). Unfiltered seawater was pumped from 2 m offshore of the adjacent reef. Flow rate through the mesocosms was maintained at a rate of 8 L min⁻¹ resulting in a seawater turnover rate of 45 min.

Experimental design

The biological response (calcification and partial mortality) of five common Hawaiian coral species (*P. compressa*, *M. capitata*, *F. scutaria*, *L. purpurea*, and *P. damicornis*) to acidification and temperature acting alone and in unison was examined over an 8-week (56-d) experimental period. Adult colony fragments of comparable size and morphology within each species were collected from the reef flat surrounding Moku o Lo'e at a depth of ~1 m. Corals were labeled for identification and 20 colonies of each species (<8 cm) were randomly assigned to one of the four experimental treatments. The experimental units were pseudoreplicated within each tank (Hurlbert 1984), but colonies of all five species

Table 1 Environmental and chemical conditions for each treatment during the 56-d experimental period

Treatment	Temperature (°C)			pCO ₂ (μatm)	DO (%)	pH _T	TA (μmol kg ⁻¹)
	Mean ± SE	Minimum	Maximum				
Ambient	27.4 ± 0.01	25.7	29.9	460 ± 13	108.9 ± 1	8.01 ± 0.01	2181 ± 9
Acidified	27.4 ± 0.01	25.7	29.6	990 ± 86	106.9 ± 2	7.74 ± 0.03	2180 ± 11
Heated	29.7 ± 0.01	26.4	31.6	475 ± 12	109.9 ± 1	8.04 ± 0.01	2191 ± 9
Acidified heated	29.5 ± 0.01	26.4	31.8	1000 ± 76	111.9 ± 1	7.72 ± 0.03	2168 ± 9

Fig. 1 Hourly irradiance levels measured continuously by cosine-corrected quantum sensor over a 56-d experimental period. Corals received maximum irradiance levels of 1914 μmol photon m⁻² s⁻¹ and mean net daily irradiance fluxes of 375.8 ± 8.8 mol photons m⁻² d⁻¹ (see Table 2)**Table 2** Irradiance characteristics for the 56-d experimental period

Maximum flux (μmol photons m ⁻² s ⁻¹)	Maximum daily flux (mol photons m ⁻² d ⁻¹)	Minimum daily flux (mol photons m ⁻² d ⁻¹)	Mean daily flux (mol photons m ⁻² d ⁻¹)
1914.0	498.5	176.1	375.8 ± 8.8

were randomly placed and each colony experienced the same conditions. The individual corals were functionally independent because the aggregated biomass (1 kg) was relatively small compared to the volume of seawater (500 L) and high turnover rates (8 L s⁻¹). We embrace the argument presented by Putnam and Edmunds (2011) that the coral fragments were not sufficiently large to change the water chemistry. In any event, all of the colonies within a treatment experienced the same physical, chemical, and biological conditions, so comparisons between species are valid.

The experimental treatments consisted of two pCO₂ levels: present-day Kāneʻohe Bay levels of 460 ± 13 μatm (mean ± SE) and double the present-day levels (990 ± 85 μatm), crossed with two temperature regimes of summer ambient 27.4 ± 0.01 °C and heated to 29.5 ± 0.01 °C (Table 1).

Chemical analyses

Midday water parameters [pH_{NBS}, salinity (‰), dissolved oxygen (% saturation), and temperature (°C)] were

measured daily to confirm desired environmental and chemical conditions (Table 1). Measurements of pH_{NBS} were taken daily using an Accumet AP72 pH/mV/temperature meter (accuracy ±0.01 pH and ±0.02 mV) and confirmed spectrophotometrically twice a week using *m*-cresol purple dye (Sigma-Aldrich #857890) according to SOP 7 (Dickson et al. 2007). Total alkalinity (TA) was measured independently twice a week using an automatic titrator (Titrino Plus 877, Metrohm) with pH glass electrode (9101 Herisau, Metrohm) to verify the accuracy and precision of the pCO₂ manipulation mechanism (Jokiel et al. 2014a). Desired precision of the automatic titrator was confirmed with certified reference materials (Batch 127 from A. Dickson Laboratory, Scripps Institution of Oceanography) twice a week. Temperature, TA, pH_{Total}, and salinity were used to calculate pCO₂ in CO2SYS (Pierrot et al. 2006) with the stoichiometric dissociation constants (K₁, K₂) defined by Mehrbach et al. (1973) and refit by Dickson and Millero (1987). Daily irradiance reaching the corals within the mesocosms was measured using a cosine-corrected quantum sensor (LI-250A meter,




Fig. 2 a Temperature ($^{\circ}\text{C}$) and b pH_{NBS} as a function of irradiance ($\mu\text{mol photon m}^{-2} \text{s}^{-1}$) in each experimental mesocosm: ambient (open squares), acidified (solid squares), heated (open circles), and acidified heated (solid circles)

LI-7792 sensor). Onset Hobo temperature loggers (UA-001-64, accuracy ± 0.53 $^{\circ}\text{C}$) were placed into each treatment, recording at 10-min intervals for the duration of the experiment. Additionally, diurnal variability in environmental parameters [pH_{NBS} , salinity (‰), dissolved oxygen (% saturation), and temperature ($^{\circ}\text{C}$)] among treatments were confirmed by placing a YSI sonde (model 6920V2-S) in each mesocosm for a 24-h period programmed at 5-min intervals (Fig. 2; Electronic Supplementary Material, ESM, Table S1).

Calcification

Coral calcification rates (mm d^{-1}) were measured throughout the 56-d experimental period using the buoyant weight method (Jokiel et al. 1978). This technique is a nondestructive and sensitive measurement of the skeletal mass, which allows for multiple measurements over time. Corals ($n = 400$) were weighed every 2 weeks for 56 d. The initial and final buoyant weights were converted to dry skeletal weight for each species (Jokiel et al. 1978). These data were expressed as the mean solid radius, which uses cube-root approximation to compute a one-dimension linear estimate (Maragos 1978). Therefore, the calcification

rate is expressed as a change in length of the radius rather than the weight change. This transformation permits comparison of colony calcification independent of coral sizes and morphology (Maragos 1978).

Partial mortality

Partial mortality was defined as percentage of dead skeletal area on each coral colony. Partial mortality was scored in bins of 5 % twice per week. Values ranged from zero (no mortality) through various amounts of tissue loss (partial mortality) to 100 % (whole-colony mortality) (Baird and Marshall 2002). To reduce observer variability, two experienced investigators conducted repeated measures until estimated tissue loss was < 5 %.

Statistical treatment of data

Coral calcification and partial mortality were measured throughout the 56-d experimental period. Mortality began after 42 d of exposure to the experimental treatments and proceeded rapidly. Therefore, coral calcification rates (mm d^{-1}) were analyzed at 42 d (no mortality) and at the end of the experiment (56 d). A two-way ANOVA model with fixed factors of pCO_2 and temperature by species was used to analyze the 42-d calcification data. Coral calcification at the end of the experiment was analyzed using a general linear model (GLM). Corals with high levels of partial mortality (> 50 % tissue loss) at the end of the 56-d experiment were removed from the GLM calcification analysis. Type III sums of squares were used to estimate the main effect of the squared differences of the unweighted marginal means.

Mortality was analyzed at the end of the full 8-week experimental stress period. Because data were percentages, partial mortality data were adjusted using an arcsine cube-root transformation and subsequently analyzed using a two-way ANOVA model with fixed factors of temperature and pCO_2 by species. Assumptions of normal distribution and homoscedasticity were assessed through graphical analyses of the residuals. Descriptive and statistical analyses were conducted using JMP Pro 11 (SAS Institute Inc., USA).

Among the response variables, interactions between stressors were described as additive, synergistic, or antagonistic within each species (Schmidt et al. 2014). If both individual factors were significant and the interaction was not, the combined effects of the stressors were equal to the sum of the individual effects; therefore, the interaction was considered additive. In cases where the statistical model indicated a significant interaction between the stressors, the combined effects were assessed by the percentage change between the responses of the corals in the treatment (least squared means) compared with the response of the corals in

the control (27.4 °C, 460 μatm) within each species. If the combined effect exceeded the sum of the individual effects, the interaction was considered to be synergistic. Conversely, if the combined effect of the stressors was less than the sum of the individual effects, the interaction was considered to be antagonistic (Table 3).

Results

Treatment conditions

During the experiment, the corals received full natural solar radiation at midday maximum irradiance levels of 1914 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ and mean net daily irradiance fluxes of $375.8 \pm 8.8 \text{ mol photons m}^{-2} \text{d}^{-1}$ (Fig. 1; Table 2). Manipulative treatments of temperature (one-way ANOVA; $F_{(3,187)} = 100.35$; $p = 0.6660$) and acidification (one-way ANOVA; $F_{(3,43)} = 34.02$; $p = 0.4746$) were not significantly different among mesocosms.

Calcification

Compared to control treatment (27.4 °C), increased temperatures (29.5 °C) significantly reduced calcification rates (pooled among pCO_2 ; $n = 80$, mean difference \pm SE) in *P. damicornis* (−58 %; $-0.20 \pm 0.05 \text{ mm d}^{-1}$; $p < 0.0001$), *M. capitata* (−55 %; $-0.15 \pm 0.05 \text{ mm d}^{-1}$; $p < 0.0001$), and *P. compressa* (−89 %; $-0.27 \pm 0.05 \text{ mm d}^{-1}$; $p < 0.0001$) after 42 d of exposure (Fig. 3;

ESM Table S2). Elevated temperatures did not have a significant effect on calcification of *F. scutaria* ($p = 0.108$) or *L. purpurea* ($p = 0.7114$). Increased pCO_2 to double the present-day levels significantly reduced calcification (pooled among temperature; $n = 80$) in *P. damicornis* (−28 %; $-0.08 \pm 0.03 \text{ mm d}^{-1}$; $p = 0.019$) and *P. compressa* (−92 %; $-0.28 \pm 0.05 \text{ mm d}^{-1}$; $p < 0.0001$) when compared to control corals (Table 3). The calcification rates of *F. scutaria* ($p = 0.070$), *L. purpurea* ($p = 0.279$), and *M. capitata* ($p = 0.056$) were not significantly suppressed under acidified conditions. Elevated temperatures and increased pCO_2 revealed an antagonistic interaction on the calcification response of *P. damicornis* with an average decline of 66 % ($-0.28 \pm 0.03 \text{ mm d}^{-1}$) (Table 3). Temperature and pCO_2 did not have a combined effect on the other tested species (Fig. 2; Table 3).


Over the 56-d exposure to experimental treatments, elevated temperature significantly reduced calcification rates (pooled among pCO_2) in *P. damicornis* (−69 %; $-0.022 \pm 0.003 \text{ mm d}^{-1}$; $n = 63$; $p < 0.0001$), *F. scutaria* (−23.9 %; $-0.015 \pm 0.004 \text{ mm d}^{-1}$; $n = 80$; $p = 0.0391$), and *M. capitata* (−67.4 %; $-0.019 \pm 0.002 \text{ mm d}^{-1}$; $n = 64$; $p < 0.0001$) (ESM Fig. S1; ESM Table S2). Increased pCO_2 significantly reduced calcification (pooled among temperature) in *P. damicornis* (−38.6 %; $-0.009 \pm 0.003 \text{ mm d}^{-1}$; $n = 63$; $p = 0.0372$) and *F. scutaria* (−26.4 %; $-0.013 \pm 0.004 \text{ mm d}^{-1}$; $n = 80$; $p = 0.0196$) (ESM Fig. S1; ESM Table S2). Statistical analysis of the 56-d calcification period revealed no significant interactions between elevated temperature and pCO_2 among tested species.

Table 3 Summary of the individual and combined effects of increased temperature and acidification stress on the biological responses of the tested coral species based on the statistical models given in ESM Table S1

Biological response	Summary of linear model results			Fractional changes compared to control			Additive interaction prediction	
	Temp	pCO_2	Temp \times pCO_2	Temp	pCO_2	Temp \times pCO_2	A + B − (A \times B)	Combined effect
Calcification (mm d^{-1}) 42 d								
<i>Pocillopora damicornis</i>	<0.0001	0.019	0.029	−0.576	−0.289	−0.662	−1.031	Antagonistic
<i>Fungia scutaria</i>	0.180	0.070	0.293	n.a.	n.a.	n.a.	n.a.	n.a.
<i>Leptastrea purpurea</i>	0.416	0.279	0.073	n.a.	n.a.	n.a.	n.a.	n.a.
<i>Montipora capitata</i>	<0.0001	0.056	0.095	−0.549	n.a.	n.a.	n.a.	Only temp effect
<i>Porites compressa</i>	<0.0001	<0.0001	0.196	−0.892	−0.919	n.a.	n.a.	Additive
Partial Mortality (% dead skeletal area)								
<i>Pocillopora damicornis</i>	<0.0001	0.752	0.058	1.825	n.a.	n.a.	n.a.	Only temp effect
<i>Montipora capitata</i>	<0.0001	0.118	0.283	0.675	n.a.	n.a.	n.a.	Only temp effect
<i>Porites compressa</i>	<0.0001	<0.0001	0.012	0.649	0.715	2.543	0.900	Synergy

Interpretation of combined effects is based on calculated predicted additive inhibition based on the fractional changes of treated corals compared with control corals. N.A. indicates that the single factor and/or interaction terms are not significant. Significant p values ($p < 0.05$) are shown in bold type. There were no significant interactions between elevated temperature and pCO_2 among tested species, except in partial mortality of *P. compressa*

Fig. 3 Calcification (mm d^{-1}) of **a** *Pocillopora damicornis*, **b** *Montipora capitata*, **c** *Porites compressa*, **d** *Fungia scutaria*, and **e** *Leptastrea purpurea* after 42-d exposure to ambient (ambient, 27.4 ± 0.01 °C) and elevated temperatures (heated, 29.5 ± 0.01 °C) and ambient (black bars; pH 8.02 ± 0.01) and high pCO_2 (acidified; blue bars; pH 7.75 ± 0.03) conditions. Error bars represent standard error of mean ($n = 20$)



Partial mortality

At the end of the 56-d experimental period, mortality was not observed in *F. scutaria* ($n = 80$) or *L. purpurea* ($n = 80$). Elevated temperature caused high rates of tissue loss leading to significant increases in exposed dead skeleton (pooled among pCO_2 ; $n = 80$) in *P. damicornis* (183 %), *M. capitata* (67 %), and *P. compressa* (65 %) (Fig. 3; Table 3). Elevated pCO_2 (pooled among temperature; $n = 80$) increased dead skeletal area of *P. compressa* (71 %) (Table 3). Temperature and pCO_2 interacted synergistically in *P. compressa* ($p = 0.012$), increasing dead skeletal area by over 254 % in comparison with control corals (Fig. 4; Table 3). Elevated temperature and pCO_2 did not have a combined effect on the other species.

Discussion

Results of this experiment support observations that biological response of calcifying organisms to temperature and pCO_2 elevation is highly variable and species-specific (Rodolfo-Metalpa et al. 2010; Erez et al. 2011; Edmunds et al. 2012; Comeau et al. 2013b; Schmidt et al. 2014). In this study, increased temperature was the dominant factor controlling reef coral calcification and mortality, with

increased pCO_2 playing a lesser role. Coral species' responses to pCO_2 and elevated temperatures acting independently and in unison varied across tested species. The synergistic effect between high temperature and high pCO_2 was weak or nonexistent among the five species tested. The interaction of elevated temperature and increased acidification showed a reduced effect on calcification rates of *P. damicornis* suggesting that acidification may ameliorate the effects of high temperature on this species. Conversely, the calcification rate of *M. capitata* was suppressed by high temperature and was not influenced by acidification. The individual effects of temperature and acidification suppressed calcification rates in *P. compressa*, but the effects were not further exacerbated by their interaction. Similarly, long-term exposure (56 d) to elevated temperature and acidification independently suppressed calcification rates in *F. scutaria*, but the response was not observed after 42 d of exposure. Mortality was mainly driven by high temperature, but interacted synergistically with pCO_2 in *P. compressa*. Elevated temperature and increased pCO_2 and their interaction did not influence the biological response of *L. purpurea*.

Thus, the combined effects of acidification and increased seawater temperatures revealed antagonistic, synergistic, and additive effects. A variety of interactive effects of temperature and pCO_2 have been reported for




Fig. 4 Partial mortality (percent dead tissue) of **a** *Pocillopora damicornis*, **b** *Montipora capitata*, and **c** *Porites compressa* at the end of the 56-d experiment under ambient (ambient; 27.4 ± 0.01 °C) and elevated temperatures (heated; 29.5 ± 0.01 °C) and ambient

(black bars; pH 8.02 ± 0.01) and high pCO₂ (acidified; blue bars; pH 7.75 ± 0.03) conditions. Mortality was not observed in *Fungia scutaria* or *Leptastrea purpurea*. Error bars represent standard error of mean ($n = 20$)

different species. For example, Reynaud et al. (2003) and Anlauf et al. (2011) showed negative synergistic effects of increased temperatures under acidification on *Stylophora pistillata* and *Porites panamensis*.

Increases in mortality resulted in unbalanced 56-d calcification data in *M. capitata*, *P. compressa*, and *P. damicornis*. This may lead to multicollinearity, therefore reducing the power of the design and making it possible to overlook the effects of the factors (Shaw and Mitchell-Olds 1993). Thus, we have focused on the interpretation of the 42-d calcification analyses for these species. No mortality was observed in *L. purpurea* or *F. scutaria*. Elevated temperature and acidification had no influence on the calcification rates of *L. purpurea* regardless of the length of exposure. Conversely, reductions in calcification were observed in *F. scutaria* after long-term exposure to elevated temperature and pCO₂.

The species responses observed in this study and others can be attributed to differences in properties of both the coral host and symbiont (Baird et al. 2009). These differences may include holobiont metabolism, heterotrophic carbon contribution capability (Grottoli et al. 2006), tissue composition and thickness (Hoegh-Guldberg 1999; Loya et al. 2001), mass-transfer rates (Nakamura and Van Woesik 2001), physiological and photosynthetic differences among the symbionts (Rowan and Powers 1991; Iglesias-Prieto and Trench 1994; Warner et al. 1996; Stimson et al. 2002; Baker 2003), species morphology (Loya et al. 2001), or other environmental and physiological conditions.

Branched corals appear to be more susceptible to thermal stress than massive and encrusting forms (Loya et al. 2001). For example, the hardy encrusting coral, *L. purpurea*, is found in extreme environments (e.g., tide pools, ship harbors, reef flats) (Coles et al. 1999; Bahr et al. 2015a) and has been shown to survive and increase in

relative abundance during reoccurring thermal events (Loya et al. 2001; van Woesik et al. 2013). In addition, *P. damicornis* was susceptible to previous bleaching events in Hawai'i (Jokiel and Brown 2004; Bahr et al. 2015b) and prolonged exposure to high temperatures led to high mortality levels within 2 weeks (Mayfield et al. 2013). Elevated temperatures decrease the ratio of photosynthesis to respiration (Coles and Jokiel 1977). *Fungia scutaria* has higher ratios of photosynthesis to respiration, higher lethal temperatures, and higher rates of heterotrophy than *M. capitata* and *P. damicornis* (Coles and Jokiel 1977). *Montipora capitata* and *P. compressa* exhibited different lipid compositions and lipid depletion rates following a bleaching event (Grottoli et al. 2006). *Montipora capitata* was better able to support daily metabolic energy requirements by its relatively low respiration rate and heterotrophic carbon contribution capability in comparison with *P. compressa*. Additionally, *F. scutaria* completely bleached during this experiment and was only significantly affected after long-term exposure to elevated temperature or increase pCO₂. Recovery of *F. scutaria* was observed within 60 d after the stressors were removed. Therefore, in corals with low respiration rates and high heterotrophic capability, lipid reserves may be able to offset energetic costs of high temperature.

High pCO₂ may reduce calcification through increased metabolic cost and photochemical impairment (Edmunds 2012). Therefore, faster growing corals may be more vulnerable to acidification (Jokiel 2011; Comeau et al. 2013a; Jokiel et al. 2014b). The faster growing corals (*P. damicornis*, *P. compressa*) in this experiment were more susceptible to acidification stress and temperature stress. *Montipora capitata*, *F. scutaria*, and *L. purpurea* showed low sensitivity to high pCO₂, which agrees with results reported for other species including additionally fed massive *Porites* spp. (Edmunds 2012), the temperate coral

Cladocora caespitosa (Rodolfo-Metalpa et al. 2010), and *Stylophora pistillata* (Reynaud et al. 2003). Lack of sensitivity in a species may be attributed to higher heterotrophic capability, lower respiration rates, and lower calcification rates, which are more dependent on temperature. Coral species with these attributes (high rates of heterotrophy and lower respiration rates) may have an ecological advantage over other species under these conditions. Coral populations that have the ability to feed heterotrophically, compile large lipid reserves, and reduce their metabolic rate may display high resistance and resilience to climate change stressors (Grottoli et al. 2006; Cohen and Holcomb 2009). This may lead to a species composition shift on future reefs (Grottoli et al. 2006).

Variation in the biological response to temperature and pCO₂ stressors may lie in regulation of material fluxes between the coral boundary layers as well as energy allocation and storage. Environmental variability may also influence conflicting results among experiments. Irradiance has been shown to interact with temperature as well as pCO₂ to produce variability in the biological response of corals (Jokiel and Coles 1977; Dufault et al. 2013; Suggett et al. 2013). Therefore, further experimentation is needed to test the interactive effects of temperature, pCO₂, and irradiance in combination. Conclusions on the biological response of corals to future climate change conditions drawn from manipulative experiments with unnatural physical and chemical water conditions may be misleading (Edmunds et al. 2012). It is therefore important to consider the extent of the experimental manipulations, degree of manipulation, the duration of the experiment, and irradiance regimes in biological responses to climate change stressors.

In general, increased seawater temperatures played the dominant role in influencing coral calcification and mortality among the tested Hawaiian species. The overall response of this five-species mixed coral community was negative as shown by decreased calcification and increased mortality among many of the species. Biological response to temperature and pCO₂ elevation is highly species-specific, so care must be taken when applying generalizations based on response of a single species to diverse and complex coral reef communities.

Acknowledgments This study was funded by the Pacific Island Climate Change Cooperative agreement number 12170-B-G101, the American Museum of Natural History Lerner Gray Memorial Fund, and the University of Hawai'i Charles and Margaret Edmondson Research Fund. We are grateful for the laboratory assistance from A. Peterson, L. Rice, M. Kosma, and F. Farrell as well as statistical guidance from A. Taylor. We would like to thank four anonymous reviewers for their time, advice, and comments which have helped improve the manuscript significantly. This is the Hawai'i Institute of Marine Biology (HIMB) contribution #1645 and the School of Ocean and Earth Science and Technology (SOEST) contribution #9558.

Corals were collected under the Hawai'i Institute of Marine Biology Special Activity Permit 2013-7.

References

- Anlauf H, D'Croz L, O'Dea A (2011) A corrosive concoction: the combined effects of ocean warming and acidification on the early growth of a stony coral are multiplicative. *J Exp Biol* 397:13–20
- Bahr KD, Jokiel PL, Toonen RJ (2015a) The unnatural history of Kāne'ohe Bay: coral reef resilience in the face of centuries of anthropogenic impacts. *Peer J* 3:e950
- Bahr KD, Jokiel PL, Rodgers KS (2015b) The 2014 coral bleaching and freshwater flood events in Kāne'ohe Bay, Hawai'i. *Peer J* 3:e1136
- Baird A, Marshall P (2002) Mortality, growth and reproduction in scleractinian corals following bleaching on the Great Barrier Reef. *Mar Ecol Prog Ser* 237:133–141
- Baird AH, Bhagooli R, Ralph PJ, Takahashi S (2009) Coral bleaching: the role of the host. *Trends Ecol Evol* 24:16–20
- Baker AC (2003) Flexibility and specificity in coral-algal symbiosis: diversity, ecology, and biogeography of Symbiodinium. *Annu Rev Ecol Syst* 2003:661–689
- Ban SS, Graham NA, Connolly SR (2014) Evidence for multiple stressor interactions and effects on coral reefs. *Glob Chang Biol* 20:681–697
- Brown BE (1997) Coral bleaching: causes and consequences. *Coral Reefs* 16:S129–S138
- Buddemeier RW, Jokiel PL, Zimmerman KM, Lane DR, Carey JM, Bohling GC, Martinich JA (2008) A modeling tool to evaluate regional coral reef responses to changes in climate and ocean chemistry. *Limnol Oceanogr Methods* 6:395–411
- Cohen AL, Holcomb M (2009) Why corals care about ocean acidification: uncovering the mechanism. *Oceanography* 22:118–127
- Coles SL, Jokiel PL (1977) Effects of temperature on photosynthesis and respiration in hermatypic corals. *Mar Biol* 43:209–216
- Coles SL, DeFelice R, Eldredge L, Carlton J (1999) Historical and recent introductions of non-indigenous marine species into Pearl Harbor, Oahu, Hawaiian Islands. *Mar Biol* 135:147–158
- Comeau S, Carpenter RC, Edmunds PJ (2013a) Response to coral reef calcification: carbonate, bicarbonate and proton flux under conditions of increasing ocean acidification. *Proc Biol Sci* 280:20131153
- Comeau S, Edmunds PJ, Spindel N, Carpenter RC (2013b) The responses of eight coral reef calcifiers to increasing partial pressure of CO₂ do not exhibit a tipping point. *Limnol Oceanogr* 58:388–398
- Comeau S, Carpenter R, Lantz C, Edmunds P (2015) Ocean acidification accelerates dissolution of experimental coral reef communities. *Biogeosciences* 12:365–372
- Dickson AG, Millero FJ (1987) A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Res A* 34:1733–1743
- Dickson AG, Sabine CL, Christian JR (2007) Guide to best practices for ocean CO₂ measurements. PICES Special Publication 3, North Pacific Marine Science Organization, Sidney, British Columbia
- Dufault AM, Ninokawa A, Bramanti L, Cumbo VR, Fan TY, Edmunds PJ (2013) The role of light in mediating the effects of ocean acidification on coral calcification. *J Exp Biol* 216:1570–1577
- Edmunds PJ (2012) Effect of pCO₂ on the growth, respiration, and photophysiology of massive *Porites* spp. in Moorea, French Polynesia. *Mar Biol* 159:2149–2160

- Edmunds PJ, Brown D, Moriarty V (2012) Interactive effects of ocean acidification and temperature on two scleractinian corals from Moorea, French Polynesia. *Glob Chang Biol* 18:2173–2183
- Erez J, Reynaud S, Silverman J, Schneider K, Allemand D (2011) Coral calcification under ocean acidification and global change. In: Dubinsky Z, Stambler N (eds) *Coral reefs: an ecosystem in transition*. Springer, Amsterdam, pp 151–176
- Fabricius K, De'ath G, Noonan S, Uthicke S (2014) Ecological effects of ocean acidification and habitat complexity on reef-associated macroinvertebrate communities. *Proc Biol Sci* 281:20132479
- Gattuso J-P, Hansson L (2011) Ocean acidification: background and history. *Ocean Acidif* 2011:1–20
- Gattuso J-P, Allemand D, Frankignoulle M (1999) Photosynthesis and calcification at cellular, organismal and community levels in coral reefs: a review on interactions and control by carbonate chemistry. *Am Zool* 39:160–183
- Gattuso JP, Frankignoulle M, Bourge I, Romaine S, Buddemeier RW (1998) Effect of calcium carbonate saturation of seawater on coral calcification. *Glob Planet Change* 18:37–46
- Glynn PW, D'Croz L (1990) Experimental evidence for high temperature stress as the cause of El Niño-coincident coral mortality. *Coral Reefs* 8:181–191
- Grottoli AG, Rodrigues LJ, Palardy JE (2006) Heterotrophic plasticity and resilience in bleached corals. *Nature* 440:1186–1189
- Hoegh-Guldberg O (1999) Climate change, coral bleaching and the future of the world's coral reefs. *Mar Freshw Res* 50:839–866
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N, Bradbury RH, Dubi A, Hatzilios ME (2007) Coral reefs under rapid climate change and ocean acidification. *Science* 318:1737–1742
- Hoeke RK, Jokiel PL, Buddemeier RW, Brainard RE (2011) Projected changes to growth and mortality of Hawaiian corals over the next 100 years. *PLoS One* 6:e18038
- Hofmann GE, Barry JP, Edmunds PJ, Gates RD, Hutchins DA, Klinger T, Sewell MA (2010) The effect of ocean acidification on calcifying organisms in marine ecosystems: an organism-to-ecosystem perspective. *Annu Rev Ecol Evol Syst* 41:127–147
- Hurlbert SH (1984) Pseudoreplication and the design of ecological field experiments. *Ecol Monogr* 54:187–211
- Iglesias-Prieto R, Trench RK (1994) Acclimation and adaptation to irradiance in symbiotic dinoflagellates. I. Responses of the photosynthetic unit to changes in photon flux density. *Mar Ecol Prog Ser* 113:163–175
- Jokiel PL (2011) The reef coral two compartment proton flux model: a new approach relating tissue-level physiological processes to gross corallum morphology. *J Exp Biol* 409:1–12
- Jokiel PL, Coles SL (1977) Effects of temperature on the mortality and growth of Hawaiian reef corals. *Mar Biol* 43:201–208
- Jokiel PL, Coles SL (1990) Response of Hawaiian and other Indo-Pacific reef corals to elevated temperature. *Coral Reefs* 8:155–162
- Jokiel PL, Brown EK (2004) Global warming, regional trends and inshore environmental conditions influence coral bleaching in Hawaii. *Glob Chang Biol* 10:1627–1641
- Jokiel PL, Maragos JE, Franzisket L (1978) Coral growth: buoyant weight technique. In: Stoddart DR, Johannes RE (eds) *Coral reefs: research methods*. Monographs on oceanographic methodology, UNESCO, Paris, pp 529–542
- Jokiel PL, Bahr KD, Rodgers KS (2014a) Low cost ocean acidification system for high-flow mesocosm and field applications on coral reefs. *Limnol Oceanogr Methods* 12:313–322
- Jokiel PL, Jury CP, Rodgers KS (2014b) Coral-algae metabolism and diurnal changes in the CO₂-carbonate system of bulk sea water. *Peer J* 2:e378
- Jokiel PL, Rodgers KS, Kuffner IB, Andersson AJ, Cox EF, Mackenzie FT (2008) Ocean acidification and calcifying reef organisms: a mesocosm investigation. *Coral Reefs* 27:473–483
- Kleypas JA, Langdon C (2006) Coral reefs and changing seawater carbonate chemistry. In: Phinney JT, Hoegh-Guldberg O, Kleypas J, Skirving W, Strong A (eds) *Coastal and estuarine studies: coral reefs and climate change science and management*. American Geophysical Union, Washington D.C., pp 73–110
- Kleypas JA, Buddemeier RW, Archer D, Gattuso JP, Langdon C, Opdyke BN (1999) Geochemical consequences of increased atmospheric carbon dioxide on coral reefs. *Science* 284:118–120
- Kleypas JA, Feely RA, Fabry VJ, Langdon C, Sabine CL, Robbins LL (2006) Impacts of ocean acidification on coral reefs and other marine calcifiers: a guide for future research. In: Report of a workshop sponsored by NSF, NOAA, and the US Geological Survey, St Petersburg, FL
- Kroeker KJ, Kordas RL, Crim R, Hendriks IE, Ramajo L, Singh GS, Duarte CM, Gattuso JP (2013) Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Glob Chang Biol* 19:1884–1896
- Lacis AA, Schmidt GA, Rind D, Ruedy RA (2010) Atmospheric CO₂: principal control knob governing Earth's temperature. *Science* 330:356–359
- Langdon C, Atkinson M (2005) Effect of elevated pCO₂ on photosynthesis and calcification of corals and interactions with seasonal change in temperature/irradiance and nutrient enrichment. *J Geophys Res Oceans* 110:C09S07
- Langdon C, Takahashi T, Sweeney C, Chipman D, Goddard J, Marubini F, Aceves H, Barnett H, Atkinson MJ (2000) Effect of calcium carbonate saturation state on the calcification rate of an experimental coral reef. *Global Biogeochem Cycles* 14:639–654
- Lesser MP, Stochaj WR, Tapley DW, Chick JM (1990) Bleaching in coral reef anthozoans: effects of irradiance, ultraviolet radiation, and temperature on the activities of protective enzymes against active oxygen. *Coral Reefs* 4:225–232
- Loya Y, Sakai K, Yamazato K, Nakano Y, Sambali H, van Woesik R (2001) Coral bleaching: the winners and the losers. *Ecol Lett* 4:122–131
- Maragos J (1978) Coral growth: geometrical relationships. In: Stoddart DR, Johannes RE (eds) *Handbook of coral reef research methods*. UNESCO, Paris, pp 543–550
- Marubini F, Ferrier-Pages C, Cuif JP (2003) Suppression of skeletal growth in scleractinian corals by decreasing ambient carbonate-ion concentration: a cross-family comparison. *Proc Biol Sci* 270:179–184
- Marubini F, Barnett H, Langdon C, Atkinson MJ (2001) Dependence of calcification on light and carbonate ion concentration for the hermatypic coral *Porites compressa*. *Mar Ecol Prog Ser* 220:153–162
- Mayfield AB, Chen M-N, Meng P-J, Lin H-J, Chen C-S, Liu P-J (2013) The physiological response of the reef coral *Pocillopora damicornis* to elevated temperature: results from coral reef mesocosm experiments in Southern Taiwan. *Mar Environ Res* 86:1–11
- Mehrbach C, Culbertson CH, Hawley JE (1973) Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnol Oceanogr* 18:897–907
- Müller A, Gagan MK, Lough JM (2004) Effect of early marine diagenesis on coral reconstructions of surface-ocean 13C/12C and carbonate saturation state. *Global Biogeochem Cycles* 18:GB1033
- Nakamura T, Van Woesik R (2001) Water-flow rates and passive diffusion partially explain differential survival of corals during the 1998 bleaching event. *Mar Ecol Prog Ser* 212:301–304

- Pandolfi JM, Connolly SR, Marshall DJ, Cohen AL (2011) Projecting coral reef futures under global warming and ocean acidification. *Science* 333:418–422
- Pierrot D, Lewis E, Wallace DWR (2006) MS Excel program developed for CO₂ system calculations. ORNL/CDIAC-105a. Carbon Dioxide Information Analysis Center. Oak Ridge National Laboratory, TN
- Putnam HM, Edmunds PJ (2011) The physiological response of reef corals to diel fluctuations in seawater temperature. *J Exp Biol* 396:216–223
- Reynaud S, Leclercq N, Romaine-Lioud S, Ferrier-Pagès C, Jaubert J, Gattuso JP (2003) Interacting effects of CO₂ partial pressure and temperature on photosynthesis and calcification in a scleractinian coral. *Glob Chang Biol* 9:1660–1668
- Rodolfo-Metalpa R, Martin S, Ferrier-Pagès C, Gattuso J-P (2010) Response of the temperate coral *Cladocora caespitosa* to mid- and long-term exposure to pCO₂ and temperature levels projected for the year 2100 AD. *Biogeosciences* 7:289–300
- Rowan R, Powers DA (1991) A molecular genetic classification of zooxanthellae and the evolution of animal-algal symbioses. *Science* 251:1348–1351
- Schmidt C, Kucera M, Uthicke S (2014) Combined effects of warming and ocean acidification on coral reef Foraminifera *Marginopora vertebralis* and *Heterostegina depressa*. *Coral Reefs* 33:805–818
- Shaw RG, Mitchell-Olds T (1993) ANOVA for unbalanced data: an overview. *Ecology* 74:(6)1638–1645
- Smith SV, Buddemeier R (1992) Global change and coral reef ecosystems. *Annu Rev Ecol Syst* 1992:89–118
- Stimson J, Sakai K, Sembali H (2002) Interspecific comparison of the symbiotic relationship in corals with high and low rates of bleaching-induced mortality. *Coral Reefs* 21:409–421
- Suggett DJ, Dong LF, Lawson T, Lawrenz E, Torres L, Smith DJ (2013) Light availability determines susceptibility of reef building corals to ocean acidification. *Coral Reefs* 32:327–337
- van Woesik R, van Woesik K, van Woesik L, van Woesik S (2013) Effects of ocean acidification on the dissolution rates of reef-coral skeletons. *Peer J* 1:e208
- Warner ME, Fitt WK, Schmidt GW (1996) The effects of elevated temperature on the photosynthetic efficiency of zooxanthellae in hospite from four different species of reef coral: a novel approach. *Plant Cell Environ* 19:291–299